```
1
       10A NCAC 41A .0101 is proposed as a temporary rule as follows:
 2
 3
                                  REPORTABLE DISEASES AND CONDITIONS
       10A NCAC 41A .0101
 4
       (a) The following named diseases and conditions are declared to be dangerous to the public health and are hereby made
       reportable within the time period specified after the disease or condition is reasonably suspected to exist:
 5
                         acquired immune deficiency syndrome (AIDS) - 24 hours;
 6
                (1)
 7
                (2)
                         anthrax - immediately;
                         botulism - immediately;
 8
                (3)
 9
                (4)
                         brucellosis - 7 days;
10
                (5)
                         campylobacter infection - 24 hours;
11
                (6)
                         chancroid - 24 hours;
12
                (7)
                         chikungunya virus infection - 24 hours;
                         chlamydial infection (laboratory confirmed) - 7 days;
13
                (7)(8)
14
                <del>(8)</del>(9)
                         cholera - 24 hours;
15
                (9)(10) Creutzfeldt-Jakob disease – 7 days;
16
                \frac{(10)}{(11)} cryptosporidiosis – 24 hours;
17
                \frac{(11)}{(12)} cyclosporiasis – 24 hours;
18
                (12)(13) dengue - 7 days;
19
                (13)(14) diphtheria - 24 hours;
20
                (14)(15) Escherichia coli, shiga toxin-producing - 24 hours;
21
                (15)(16) ehrlichiosis – 7 days;
22
                (16)(17) encephalitis, arboviral - 7 days;
23
                (17)(18) foodborne disease, including Clostridium perfringens, staphylococcal, Bacillus cereus, and other and
24
                         unknown causes - 24 hours;
25
                (18)(19) gonorrhea - 24 hours;
26
                (19)(20) granuloma inguinale - 24 hours;
27
                (20)(21) Haemophilus influenzae, invasive disease - 24 hours;
28
                \frac{(21)(22)}{(21)} Hantavirus infection – 7 days;
29
                (22)(23) Hemolytic-uremic syndrome – 24 hours;
30
                (23)(24) Hemorrhagic fever virus infection – immediately;
31
                \frac{(24)}{(25)} hepatitis A - 24 hours;
32
                (25)(26) hepatitis B - 24 hours;
33
                (26)(27) hepatitis B carriage - 7 days;
34
                \frac{(27)}{(28)} hepatitis C, acute – 7 days;
35
                (28)(29) human immunodeficiency virus (HIV) infection confirmed - 24 hours;
36
                (29)(30) influenza virus infection causing death – 24 hours;
37
                (30)(31) legionellosis - 7 days;
```

```
1
                (31)(32) leprosy – 7 days;
 2
                (32)(33) leptospirosis - 7 days;
 3
                (33)(34) listeriosis – 24 hours;
 4
                (34)(35) Lyme disease - 7 days;
                (35)(36) lymphogranuloma venereum - 7 days;
 5
                (36)(37) malaria - 7 days;
 6
 7
                (37)(38) measles (rubeola) - 24 hours;
 8
                (38)(39) meningitis, pneumococcal - 7 days;
 9
                (39)(40) meningococcal disease - 24 hours;
10
                (41) Middle East respiratory syndrome (MERS) - 24 hours;
11
                (40)(42) monkeypox – 24 hours;
12
                (41)(45) mumps - 7 days;
13
                (42)(44) nongonococcal urethritis - 7 days;
14
                (43)(45) novel influenza virus infection – immediately;
15
                (44)(46) plague - immediately;
                (45)(47) paralytic poliomyelitis - 24 hours;
16
17
                (46)(48) pelvic inflammatory disease – 7 days;
18
                (47)(49) psittacosis - 7 days;
19
                (48)(50) Q fever - 7 days;
20
                (49)(51) rabies, human - 24 hours;
21
                (50)(52) Rocky Mountain spotted fever - 7 days;
22
                (51)(53) rubella - 24 hours;
23
                (52)(54) rubella congenital syndrome - 7 days;
24
                (53)(55) salmonellosis - 24 hours;
25
                (54)(56) severe acute respiratory syndrome (SARS) – 24 hours;
26
                (55)(57) shigellosis - 24 hours;
27
                (56)(58) smallpox - immediately;
28
                (57)(59) Staphylococcus aureus with reduced susceptibility to vancomycin – 24 hours;
29
                (58)(60) streptococcal infection, Group A, invasive disease - 7 days;
30
                (59)(61) syphilis - 24 hours;
31
                (60)(62) tetanus - 7 days;
32
                (61)(63) toxic shock syndrome - 7 days;
33
                (62)(64) trichinosis - 7 days;
34
                (63)(65) tuberculosis - 24 hours;
35
                (64)(66) tularemia – immediately;
36
                (65)(66) typhoid - 24 hours;
37
                (66)(67) typhoid carriage (Salmonella typhi) - 7 days;
```

1	(67)(68) typhus, epidemic (louse-borne) - 7 days;						
2	<del>(68)</del> (69	(68)(69) vaccinia – 24 hours;					
3	(69)(70) vibrio infection (other than cholera) – 24 hours;						
4	<del>(70)</del> (7	(70)(71) whooping cough – 24 hours; and					
5	<del>(71)</del> (7)	<del>(71)</del> (72) yellow fever - 7 days.					
6	(b) For purpose	(b) For purposes of reporting, "confirmed human immunodeficiency virus (HIV) infection" is defined as a positive virus					
7	culture, repeate	culture, repeatedly reactive EIA antibody test confirmed by western blot or indirect immunofluorescent antibody test,					
8	positive nucleic acid detection (NAT) test, or other confirmed testing method approved by the Director of the State						
9	Public Health Laboratory conducted on or after February 1, 1990. In selecting additional tests for approval, the Director						
10	of the State Pub	lic Health	Laboratory shall consider whether such tests have been approved by the federal Food and Drug				
11	Administration, recommended by the federal Centers for Disease Control and Prevention, and endorsed by the						
12	Association of Public Health Laboratories.						
13	(c) In addition to the laboratory reports for Mycobacterium tuberculosis, Neisseria gonorrhoeae, and syphilis specified in						
14	G.S. 130A-139, laboratories shall report:						
15	(1)	(1) Isolation or other specific identification of the following organisms or their products from human					
16		clinica	I specimens:				
17		(A)	Any hantavirus or hemorrhagic fever virus.				
18		(B)	Arthropod-borne virus (any type).				
19		(C)	Bacillus anthracis, the cause of anthrax.				
20		(D)	Bordetella pertussis, the cause of whooping cough (pertussis).				
21		(E)	Borrelia burgdorferi, the cause of Lyme disease (confirmed tests).				
22		(F)	Brucella spp., the causes of brucellosis.				
23		(G)	Campylobacter spp., the causes of campylobacteriosis.				
24		(H)	Chlamydia trachomatis, the cause of genital chlamydial infection, conjunctivitis (adult and				
25			newborn) and pneumonia of newborns.				
26		(I)	Clostridium botulinum, a cause of botulism.				
27		(J)	Clostridium tetani, the cause of tetanus.				
28		(K)	Corynebacterium diphtheriae, the cause of diphtheria.				
29		(L)	Coxiella burnetii, the cause of Q fever.				
30		(M)	Cryptosporidium parvum, the cause of human cryptosporidiosis.				
31		(N)	Cyclospora cayetanesis, the cause of cyclosporiasis.				
32		(O)	Ehrlichia spp., the causes of ehrlichiosis.				
33		(P)	Shiga toxin-producing Escherichia coli, a cause of hemorrhagic colitis, hemolytic uremic				
34			syndrome, and thrombotic thrombocytopenic purpura.				
35		(Q)	Francisella tularensis, the cause of tularemia.				
36		(R)	Hepatitis B virus or any component thereof, such as hepatitis B surface antigen.				
37		(S)	Human Immunodeficiency Virus, the cause of AIDS.				

1		(T)	Legione	lla spp., the causes of legionellosis.				
2		(U)	Leptosp	ira spp., the causes of leptospirosis.				
3		(V)	Listeria	monocytogenes, the cause of listeriosis.				
4		(W)	Middle	East respiratory syndrome virus.				
5		<del>(W)</del> (X)	Monkey	rpox.				
6		( <u>X)(Y)</u>	Mycoba	cterium leprae, the cause of leprosy.				
7		<del>(Y)</del> (Z)	Plasmoo	lium falciparum, P. malariae, P. ovale, and P. vivax, the causes of malaria in humans.				
8		(Z)(AA)	<u>)</u> Poliovir	us (any), the cause of poliomyelitis.				
9		(AA)(BB)Rabies virus.						
10		(BB)(CC)Rickettsia rickettsii, the cause of Rocky Mountain spotted fever.						
11		(CC)(DD)Rubella virus.						
12		(DD)(EE)Salmonella spp., the causes of salmonellosis.						
13		(EE)(FF)Shigella spp., the causes of shigellosis.						
14		(FF)(GG)Smallpox virus, the cause of smallpox.						
15		<del>(GG)</del> (H	<u>H)</u> Staphy	clococcus aureus with reduced susceptibility to vanomycin.				
16		(HH)(II) Trichinella spiralis, the cause of trichinosis.						
17		( <del>II)</del> (JJ)	Vaccinia	a virus.				
18		<del>(JJ)</del> (KK	<u>)</u> Vibrio s	pp., the causes of cholera and other vibrioses.				
19		(KK)(LL)Yellow fever virus.						
20		(LL)(M	<u>M)</u> Yersir	nia pestis, the cause of plague.				
21	(2)	Isolation or other specific identification of the following organisms from normally sterile human body						
22		sites:						
23		(A)	Group A	A Streptococcus pyogenes (group A streptococci).				
24		(B)	Haemop	ohilus influenzae, serotype b.				
25		(C)	Neisseri	a meningitidis, the cause of meningococcal disease.				
26	(3)	Positive serologic test results, as specified, for the following infections:						
27		(A)	Fourfold	d or greater changes or equivalent changes in serum antibody titers to:				
28			(i)	Any arthropod-borne viruses associated with meningitis or encephalitis in a human.				
29			(ii)	Any hantavirus or hemorrhagic fever virus.				
30			(iii)	Chlamydia psittaci, the cause of psittacosis.				
31			(iv)	Coxiella burnetii, the cause of Q fever.				
32			(v)	Dengue virus.				
33			(vi)	Ehrlichia spp., the causes of ehrlichiosis.				
34			(vii)	Measles (rubeola) virus.				
35			(viii)	Mumps virus.				
36			(ix)	Rickettsia rickettsii, the cause of Rocky Mountain spotted fever.				
37			(x)	Rubella virus.				

1		(xi)	Yellow fever virus.		
2		(B) The pr	resence of IgM serum antibodies to:		
3		(i)	Chlamydia psittaci.		
4		(ii)	Hepatitis A virus.		
5		(iii)	Hepatitis B virus core antigen.		
6		(iv)	Rubella virus.		
7		(v)	Rubeola (measles) virus.		
8		(vi)	Yellow fever virus.		
9	(4)	Laboratory results from tests to determine the absolute and relative counts for the T-helper (CD4)			
10		subset of lymph	nocytes and all results from tests to determine HIV viral load.		
11					
12	History Note:	Authority G.S.	130A-134; 130A-135; 130A-139; 130A-141;		